

Cherax *destructor*

Species description

The common yabby (*Cherax destructor*) is a freshwater crustacean native to southeastern Australia. Initially introduced to Europe (Spain) in 1983 for aquaculture purposes, this crayfish also gained popularity in aquariums, and was therefore commercialised through the pet trade. Although discovered in a river in France in 2019, the common yabby has not yet been observed in the wild in Belgium. Its presence in European countries' natural environments is likely the result of intentional releases of aquarium specimens and experimental aquaculture. Today, while emerging in some countries, this crayfish has the potential to become a problematic aquatic invasive species in countries where it has been introduced. It was, therefore, included in the LIFE RIPARIAS alert list. Paradoxically, the common yabby has been listed as "vulnerable" in its native range by The International Union for Conservation of Nature (IUCN) Red List. Its potential presence on the Belgian territory cannot be ruled out.



Fig 1. *Cherax destructor*. Photo: Asimakis Patitsas

Cherax destructor

Cherax destructor can colonise a wide range of aquatic habitats, including ponds, streams, rivers, irrigation channels, and swamps. Its resilient nature enables it to tolerate saline environment, poorly oxygenated waters and endure exposure to extreme conditions. This is particularly crucial as most of its native range is characterised by elevated summer temperatures and limited annual rainfall. As a highly competitive invasive species, this crayfish likely exerts wide-ranging detrimental impacts on ecosystems and biodiversity, although explicit evidence remains limited. The species' aggressive nature, coupled with characteristics such as high population densities, large body size, and high fecundity, suggests it may outcompete certain native species under favourable climatic conditions. This, in turn, can lead to changes in food webs and a decline in biological diversity. The common yabby is, however, vulnerable to the *Aphanomyces astaci*, also known as the crayfish plague, also lethal to all indigenous European crayfish species, such as the threatened noble crayfish (*Astacus astacus*). Some populations in Italy appeared to have gone locally extinct due to the deadly pathogen carried by *Procambarus clarkii*. Beyond ecological impacts, economic consequences involve management-related costs and potential damages to irrigation systems.

Biological characteristics, reproduction and spread

This large crayfish exhibits burrowing behavior to seek refuge in adverse conditions. The species can excavate complex burrows as deep as 2m, that are interconnected by tunnels, and can seal the entrance with mud during extreme conditions. Juveniles, being highly sociable, often share burrows and typically maintain physical proximity to one another. This species exhibits exceptional resistance to desiccation and displays remarkable temperature tolerance, enduring water temperatures ranging between 1°C and 35°C. Persistent *Cherax destructor* populations are also found to endure, even in regions where freezing occurs. When winter temperatures threaten their survival, some crayfish species are known to burrow into underground tunnels to withstand freezing. The common yabby's growth rate is notably also influenced by temperatures, reaching its peak between 20 and 25°C.



Fig 2. *Cherax destructor*. Photo: Xavier Vermeersch

References

- Austin, C.M. (1998) Intraspecific variation in clutch and brood size and rate of development in the yabby, *Cherax destructor* (Decapoda: Parastacidae). *Aquaculture*, 167(1), 147-159.
- Coughran, J. and Daly, G. (2012) Potential threats posed by a translocated crayfish: the case of *Cherax destructor* in coastal drainages of New South Wales, Australia. *Crustacean Research*, 7, 5-13
- Gherardi, F. and Sabapathy Allen, U. (2022) *Cherax destructor* (yabby), *CABI Compendium*. <https://www.cabidigitallibrary.org/doi/10.1079/cabicompendium.89134#sec-35> [Accessed: 4th December 2023]
- Johnson, H.T. (1979) *Reproduction, development and breeding activity in the freshwater crayfish, Cherax destructor* Clark. Master of science, University of Sydney.
- Lawrence, C.G. (2001) *Morphology and incidence of Yabby (Cherax albidus) burrows in Western Australia*. Fisheries research report. Report: 129.
- Lynas, J., Storey, A.W. and Knott, B. (2007) Aggressive interactions between three species of freshwater crayfish of the genus *Cherax* (Decapoda: Parastacidae). *Marine and Freshwater Behaviour and Physiology*, 40(2), 105-116.

Like other crayfish species, the common yabby exhibits reduced activity in colder conditions, which typically starts to diminish from 15°C, to enter a condition of partial hibernation. Individuals are mostly active during the night. Their feeding behavior is largely influenced by the level of light penetration in the water, with peak activity commonly observed shortly before dawn and immediately after dusk.

Females reach sexual maturity at a young age and engage in spawning when the water temperature exceeds 15°C. In its native range, the reproductive behaviour of *Cherax destructor* follows a distinct seasonal pattern, marked by mating and spawning events during spring. The common yabby is able to undergo multiple spawning events, occurring up to five times a year, if the right water temperatures are met (between 18°C and 20°C). Female nurtures the eggs by attaching them beneath their abdomen. The typical number of eggs in a clutch is around 400 but can reach over 1000 for a large female. The species possesses the capacity to colonise both connected and non-connected water systems. Natural overland dispersal over considerable distances, sometimes in large groups, has been observed. The species is also able to spread both downstream and upstream at important speed. Those high reproductive, adaptive, and dispersal abilities highlight the critical need for effective management measures against this species.



Fig 3. *Cherax destructor*. Photo: quollsskinks

Mazza, G. et al. (2018). The Red Alien vs. the Blue Destructor: The Eradication of *Cherax destructor* by *Procambarus clarkii* in Latium (Central Italy). *Diversity*, 10(4), 126.

Monty, A. et al. (2022) Invasive alien crayfish species - Identification and best management practices. LIFE RIPARIAS project, 50p.

Vecchioni, L., et al. (2022) An update of the known distribution and status of *Cherax* spp. in Italy (Crustacea, Parastacidae). *BiolInvasions Records*, 11(4), 1045-1055.

Verhoef, G.D. and Austin, C.M. (1999) Combined effects of temperature and density on the growth and survival of juveniles of the Australian freshwater crayfish, *Cherax destructor* Clark.; Part 1. *Aquaculture*, 170(1), 37-47.

Vesely, L., Buric, M. and Kouba, A. (2015). Hardy exotics species in temperate zone: can "warm water" crayfish invaders establish regardless of low temperatures? *Scientific Reports*, 5, 1-10.

Vigneron, T. (2019). Premier signalement de *Cherax destructor* en milieu naturel en France (Bretagne). <http://especes-exotiques-envahissantes.fr/premier-signalement-de-cherax-destructor-en-milieu-naturel-en-france-bretagne/?lang=en> [Accessed: 4th December 2023]

General considerations about management

Eradicating and controlling common yabby populations poses a considerable challenge, especially once they are well-established. To date, successful eradications have been achieved by introducing a single *Pacifastacus leniusculus* individual infected with the crayfish plague. Numerous effective management techniques, posing lower risks than introducing pathogens and diseases, are available for controlling this species. However, their successful implementation typically requires the strategic combination of methods over several years to attain satisfactory outcomes and the potential for eradication.

Given the similarity between the different crayfish species, 5 efficient methods are applicable for the management of all alien invasive crayfish species. Occasionally, species-specific information must be taken into account to allow a better application of the methods. In the case of *Cherax destructor* and its exceptional resistance to desiccation and environmental variations, the implementation of water drawdown as a management method may prove ineffective. The feasibility of eradicating common yabby populations should be evaluated on a case-by-case basis, accounting for site-specific factors and thorough discussions within the management team.

"Given the similarity between the different crayfish species, 5 efficient methods are applicable for all exotic invasive crayfish species. Occasionally, species-specific information must be taken into account to allow a better application of the methods".

Due to the species' ability to move on terrestrial and in aquatic environments between and within different water systems, precautionary measures must be implemented prior to management to prevent the dispersal of individuals to other water systems. Managed areas are, therefore, isolated by physical barriers. This precaution is especially important when the environment is disrupted by management measures, as such disturbances may heighten the likelihood of individuals seeking new favorable sites. The application of rigorous biosecurity measures, recommended to prevent the potential spread of the crayfish plague carried by certain exotic crayfish species, might be less relevant in this case. Given its vulnerability to the *Aphanomyces astaci*, the impact of the common yabby on European indigenous crayfish species such as the threatened noble crayfish (*Astacus astacus*) is probably more limited.

Managed and surrounding sites must remain under enhanced surveillance for at least 3 years after the implementation of the last treatment, and recurrent trapping implemented if necessary

Trapping

- v This method is applicable in most situations
- v Trapping has limited impact on the ecosystem
- v The method enables the monitoring of the population size (CPUE)
- x There is a risk of bycatch
- x There is a risk of exclusively capturing large males
- x The method is time consuming
- x Control can be obtained through this method. Eradication is highly unlikely
- x The real effectiveness of the method is difficult to predict

Method description

Trapping consists in placing crayfish traps in the water and collecting them at regular intervals, spanning one or several days. Ethical methods are employed to euthanise captured crayfish, minimising unnecessary suffering. The effectiveness of trapping depends on factors such as bait selection, seasonality, and weather conditions. Trapping is usually conducted in the warmer months when crayfish are most active, maximising the chances of successful captures.

This approach enables the reduction of crayfish populations, is relatively straightforward to implement, and enables the monitoring of population density through CPUE (Catch Per Unit Effort). By consistently employing the same traps and lifting them at regular intervals, the average number of captures per trap serves as an indicator of crayfish density changes over time. While trapping is a commonly employed method, accurately predicting its effectiveness poses challenges, particularly because capturing the youngest and least mobile individuals, such as ovigerous females, is complicated



Fig 4. *Procambarus clarkii* caught in a trap. Photo: Marie Patinet

References

Adriaens, T. et al. (2019). *Feasibility of eradication and spread limitation for species of Union concern sensu the EU IAS Regulation (EU 1143/2014) in Belgium*. Institute for Nature and Forest Research, Service Public de Wallonie, National Scientific Secretariat on Invasive Alien Species, Belgian Biodiversity Platform.

Basillico, L. et al. (2013). *Les invasions d'écrevisses exotiques, impacts écologiques et pistes pour la gestion*. Les rencontres de l'Onema, p. 41.

Crepin, E. (2021). *Élaboration d'un guide de bonnes pratiques pour la gestion des plantes aquatiques et des écrevisses envahissantes*, p. 223.

Gherardi, F. et al. (2011). *Managing invasive crayfish: is there a hope? Aquatic Sciences*, 73(2), 185–200.

Girardet, M.A. et al. (2012). *Eradication d'une population d'écrevisses de Louisiane, Procambarus clarkii, à l'étang de Vidy et situation des populations d'écrevisses à Lausanne, Suisse*. p. 11.

Green, N. et al. (2018). *Trapping for invasive crayfish: comparisons of efficacy and selectivity of baited traps versus novel artificial refuge traps*. *Knowledge & Management of Aquatic Ecosystems*, (419), 1-15.

To increase the chances of success, it is necessary to combine different trap designs and maintain an important capture pressure over several years. This method is time-consuming and labour-intensive, making it more suitable for small water bodies and situations where crayfish populations are still relatively limited.

When carrying out a trapping operation, it's crucial to ensure an adequate quantity of traps. Traps are available in various sizes and shapes, including cylindrical, rectangular, conical, etc. The materials commonly used are wire mesh or polyethylene netting, known for its resistance to damage caused by claws. Additionally, there are both foldable and non-foldable models.

The selected model(s) should exhibit good efficiency while minimising the risk of bycatch (such as fish, amphibians, etc.), especially in sites where these species hold high conservation value. Consistent and frequent trap collection helps mitigate mortality among unintentionally captured individuals. It's essential to disinfect and thoroughly dry traps if they are moved to a different site.

• Baited traps

Nets baited with fish, dog food, or other meat products, are commonly employed in both water bodies and slow-moving rivers. They are placed on the water bottom at a depth of at least 30-40 cm and secured to a support on the bank. It is crucial that these nets do not exclusively capture large male individuals, as this would limit or negate the impact on the population. The presence of such large individuals in a trap tends to discourage smaller ones from entering. The use of larger traps can mitigate this effect, as can restricting the size of openings: traps with larger openings (> 4 cm) predominantly capture large crayfish, whereas smaller openings (< 4 cm or even < 2 cm) facilitate the capture of smaller individuals. Additionally, the screen mesh size must be fine enough to prevent the escape of small individuals.

Holdich, D. M., Gydemo, R. and Rogers, W. D. (2017). *A review of possible methods for controlling nuisance populations of alien crayfish*. In: *Crayfish in Europe as alien species*, pp. 245-270.

Krieg, R., King, A. and Zenker, A. (2020). *Measures to Control Invasive Crayfish Species in Switzerland: A Success Story?* *Frontiers in Environmental Science*, 8, 1-9.

Poulet, N. (2014). *Les méthodes de contrôle des populations d'écrevisses invasives*, p. 13.




Fig 5. Example of baited traps Drawing : Arnaud Monty

• Fyke net

A fyke net is a very long, more or less conical net mounted on a rigid frame. It is usually equipped with side wings that guide the crayfish towards the trap entrance. Setting up a fyke net is more challenging compared to a regular net, as it needs to be anchored to the bottom with stakes, and aquatic vegetation may impede the process. In river settings, the opening is typically oriented downstream. The fyke net may be baited or left unbaited. While it has the advantage of capturing a large number of individuals and is less selective regarding crayfish size, the installation is complex, and there is a genuine risk of unintentional catches, with significant mortality.




Fig 6. Example of fyke net.
Drawing : Arnaud Monty

• Unbaited traps

Unbaited traps serve as artificial refuges, which mimic natural shelters used by crayfish such as galleries and cavities. Various structures, including PVC tubes attached to a support or bricks with holes, can be employed. The strategy involves leaving the traps in place for an extended period, typically a few days, allowing crayfish to seek refuge in the cavities before removing the traps. For optimal placement, it is recommended to place the cavities horizontally on the water body's bottom, perpendicular to the current, and secure them with stones if necessary. These traps offer the advantage of not selecting individuals based on sex or size, require no bait, and minimise risk of bycatch. Additionally, they can be used for extended periods (from April to November). However, their effectiveness, which has not been proven yet for all species, depends on the environment. While demonstrated to be effective for *Pacifastacus leniusculus* in running waters, efficacy on *Procambarus clarkii* in ponds is very low. The presence of numerous natural shelters likely diminishes their effectiveness.




Fig 7. An example of an artificial refuge trap, consisting in pieces of PVC pipe riveted onto a metal plate and blocked on one side. It may be necessary to weight this type of trap with a stone, for example

Seine fishing

Method description

- ✓ The method enables balanced catches of different sizes and sexes
- x The method demands expertise and a qualified staff for proper execution
- x The movement of operators poses potential impacts on the ecosystem.
- x While the method allows for control, achieving eradication is highly unlikely.
- x The real effectiveness of the method is difficult to predict

The seine is a large net employed by operators, typically two individuals walking in the water, that is dragged along the water's bottom during daylight hours for crayfish capture. In river settings, additional operators may flip stones and disrupt upstream vegetation to coax crayfish out of their hiding spots. Net fishing proves most effective in small, shallow water bodies that can be crossed on foot, especially those with relatively clear water.

This method facilitates the capture of juvenile crayfish and ovigerous females, offering a distinct advantage. It serves as a valuable complement to trapping, proving especially useful for harvesting significant quantities of crayfish during periods of increased activity and important population density, notably in the summer season.

However, this method does not enable local eradication and is relatively labour-intensive. The movement of operators and the overturning of obstacles constitute a non-negligible disturbance to the aquatic environment. Although the risk of bycatch is high, operators have the possibility to release individuals of non-target species.

References

Crepin, E. (2021). *Élaboration d'un guide de bonnes pratiques pour la gestion des plantes aquatiques et des écrevisses envahissantes*, p. 223.

Gherardi, F. et al. (2011). Managing invasive crayfish: is there a hope? *Aquatic Sciences*, 73(2), 185–200.

Holdich, D. M., Gydemo, R. and Rogers, W. D. (2017). A review of possible methods for controlling nuisance populations of alien crayfish. In: *Crayfish in Europe as alien species*, pp. 245-270.

Poulet, N. (2014). *Les méthodes de contrôle des populations d'écrevisses invasives*, p. 13.



Fig 8. Seine fishing

DID YOU KNOW?

Catching crayfish

Fishing for invasive alien crayfish is feasible using equipment known as "crayfish ring nets." These circular nets, equipped with ledges and attached to strings, are strategically placed on the water body's bottom. They are periodically lifted to capture crayfish. The nets are baited with pieces of fish or any other baits crayfish find palatable. String catching is another possible technique to capture crayfish.

Catching crayfish in public waters requires a permit. It is important to adhere to the specific regional regulations and designated opening periods. On private properties, the owner's authorisation is required. Although catching crayfish can be a recreational activity which may present a culinary interest, its contribution to the regulation of invasive alien crayfish is limited. On the contrary, transporting living individuals poses an increased risk of dispersal.

It is crucial to highlight that the transport of EU-listed species is strictly prohibited, with an exception allowing transportation for eradication measures. Furthermore, consuming crayfish is not recommended, as they have the potential to accumulate heavy metals or other toxic substances from polluted waters. Importantly, using invasive alien crayfish as bait for fishing is strictly forbidden.

References

Holdich, D. M., Gydemo, R. and Rogers, W. D. (2017). A review of possible methods for controlling nuisance populations of alien crayfish. In: Crayfish in Europe as alien species, pp. 245-270.



Fig 9. Crayfish ring net. Photo: Hari Seldon

Drainage

Method description

- v This method has the potential to achieve local eradication.
- v This action can be integrated in the regular management of the water body (reduction of siltation)
- x This method can have significant impacts on the ecosystem and aquatic species
- x Drainage can be an expensive method
- x The success of this approach depends on the drainage possibilities of the water body

This method involves the complete drainage of a water body to make the habitat inhospitable to crayfish, while inducing maximum mortality through additional actions. Sustained drainage over several years creates conditions that result in crayfish mortality due to desiccation or predation. A recommended duration is a 3-year drainage, spanning at least 2 winters, as crayfish exhibit resilience to both desiccation and frost.

Crayfish must be effectively contained within the desiccated area; thus, the placement of physical barriers, both on land and in the water, is essential. In the case of pumping, the discharged water must undergo filtration (using a plankton filter with a mesh size of 1 millimeter) to prevent the downstream dispersion of juveniles. Following the drainage, manual harvesting can be implemented in the subsequent days to eliminate a significant number of individuals and minimize the risk of accidental dispersion by predators. Subsequently, the application of slaked lime (calcium hydroxide) in the remaining wet areas and at the entrance of the galleries will help eliminate surviving individuals. Ideally, the drainage process should be leveraged to clean the pond's bottom and/or embank the shores, effectively destroying remaining galleries where individuals may persist. Consideration may be given to occasional concreting in high-risk areas, such as rockfill locations where individuals persist.

References

- Basilico, L. et al. (2013). *Les invasions d'écrevisses exotiques, impacts écologiques et pistes pour la gestion*. Les rencontres de l'Onema, p. 41.
- Crepin, E. (2021). *Élaboration d'un guide de bonnes pratiques pour la gestion des plantes aquatiques et des écrevisses envahissantes*, p. 223.
- Girardet, M.A. et al. (2012). *Eradication d'une population d'écrevisses de Louisiane, Procambarus clarkii, à l'étang de Vidy et situation des populations d'écrevisses à Lausanne, Suisse*. p. 11.
- Holdich, D. M., Gydemo, R. and Rogers, W. D. (2017). *A review of possible methods for controlling nuisance populations of alien crayfish*. In: *Crayfish in Europe as alien species*, pp. 245-270.
- Kouba, A. et al. (2016). *The significance of droughts for hyporheic dwellers: evidence from freshwater crayfish*. *Scientific Reports*, 6(1), 265-269.
- Krieg, R., King, A. and Zenker, A. (2020). *Measures to Control Invasive Crayfish Species in Switzerland: A Success Story?*. *Frontiers in Environmental Science*, 8, 1-9.
- Poulet, N. (2014). *Les méthodes de contrôle des populations d'écrevisses invasives*, p. 13.

If properly done, this method can enable the local eradication of a crayfish population. This has been demonstrated across various species, including *Procambarus clarkii*, which is known to be difficult to eradicate. It is crucial to emphasise that the mere implementation of routine maintenance dredging does not yield the same outcomes, and that no method can guarantee absolute eradication success.

This method has strong impacts on the aquatic environment, particularly in the medium term (over a few years). The potential transfer of non-target species to alternative sites poses risks, potentially leading to the dispersal of juvenile crayfish. Moreover, the method is associated with substantial costs and proves burdensome for users of the water body.

The method is particularly relevant for drainable water bodies of relatively modest size, where emerging populations are likely to rapidly colonize other sites.



Fig 10. Example of pond drawdown. Achieving a thorough drainage of a water body can be challenging, potentially impacting the effectiveness of the management measure. To address this challenge, various techniques can be employed, such as liming remaining water puddles or manually removing crayfish. Upper photo: Marie Patinet, lower photo: jacki-dee.



Reinforcing predator populations

Method description

- ✓ Method that can be implemented in many situations
- ✓ Limited impact on the ecosystem
- ✓ Limited costs
- ✓ Method that is easy to combine with other methods
- ✗ Difficulty in supplying some predatory species
- ✗ Method that enables control. Eradication is highly unlikely
- ✗ The real effectiveness of the method is difficult to predict

Eel (*Anguilla anguilla*), burbot (*Lota lota*), perch (*Perca fluviatilis*), and pike (*Esox lucius*) represent four native species that regularly prey on crayfish if present in the environment. Additionally, other exotic carnivorous fish, such as pike-perch (*Sander lucioperca*) or wels catfish (*Silurus glanis*), also include crayfish in their diet. Juveniles may be consumed by various species, including omnivorous fish like carp (*Cyprinus carpio*) or tench (*Tinca tinca*). Increasing predator density not only reduces crayfish populations but also limits their activity, and therefore their impact. This is a cost-effective and sustainable method. It is recommended to introduce individuals of indigenous species already present in the environment, or to encourage the growth of their populations by adjustments (water purification, spawning grounds, renaturation of banks) or regulations (protection of species, fishing restriction).

The eel is considered the most effective species in controlling crayfish populations, particularly due to its ability to consume young individuals, detect crayfish by smell, and access galleries. However, challenges in managing its reproduction and its critically endangered status present significant constraints for its use. The introduction of eels into an environment should primarily be part of a conservation plan for the safeguard of this species. If strong measures were to be taken in the future to conserve this species and restore its populations, this would probably have a positive impact on the control of alien crayfish populations.

The perch is a commonly found, sometimes abundant, small predator. Its potential impact on the youngest crayfish is particularly noteworthy, especially when combined with trapping. Although non-native, pike-perch can have a similar effect. Pike, being important consumers of crayfish, exhibit a size-dependent prey preference. Young pike feed on crayfish from their early stages, while adult individuals tend to select larger prey.

References

Adriaens, T. et al. (2019). *Feasibility of eradication and spread limitation for species of Union concern sensu the EU IAS Regulation (EU 1143/2014) in Belgium*. Institute for Nature and Forest Research, Service Public de Wallonie, National Scientific Secretariat on Invasive Alien Species, Belgian Biodiversity Platform.

Crepin, E. (2021). *Élaboration d'un guide de bonnes pratiques pour la gestion des plantes aquatiques et des écrevisses envahissantes*, p. 223.

Holdich, D. M., Gydemo, R. and Rogers, W. D. (2017). *A review of possible methods for controlling nuisance populations of alien crayfish*. In: *Crayfish in Europe as alien species*, pp. 245-270.

Krieg, R., King, A. and Zenker, A. (2020). *Measures to Control Invasive Crayfish Species in Switzerland: A Success Story?*. *Frontiers in Environmental Science*, 8, 1-9.

Poulet, N. (2014). *Les méthodes de contrôle des populations d'écrevisses invasives*, p. 13.

Various insects, such as dragonfly larvae and dytids, contribute to crayfish predation, along with birds and mammals. Beyond the occasional introduction of fish, the conservation of aquatic environments fosters a diverse predatory fauna, enhancing its ability to curb the demographic expansion of invasive alien crayfish.



Fig 11. Perch (*Perca fluviatilis*). Photo: Christa Rohrbach



Fig 12. Pike (*Esox lucius*). Photo: Gilles San Martin



Fig 13. Burbot (*Lota lota*). Photo: paul_sk11



Fig 14. Eel (*Anguilla anguilla*). Photo: Frederic-andre

Ecosystem modifications

Method description

- ✓ This approach can have positive impact on the ecosystem
- ✓ It is a long term approach
- ✗ The method requires a good knowledge of the species being managed
- ✗ The method enables control. Eradication is however highly unlikely

One of the factors contributing to the rapid formation of large populations of invasive alien crayfish is the simplification of aquatic ecosystems. Aquatic systems are increasingly becoming artificial, with features such as loose, vertical, or steep banks known to favor the construction of galleries (e.g. red swamp crayfish). These galleries, crucial for their life cycle, enable them to withstand adverse conditions like drought or frost, posing challenges for effective management. Moreover, the lack of vegetation along the banks can facilitate the dispersal of individuals and the colonisation of new water bodies. The decrease in natural predators is another element that make degraded aquatic environments particularly susceptible to invasion.

These diverse findings strongly indicate that restoring natural and diverse aquatic environments is a promising strategy for long-term control of invasions by alien crayfish. Implementing measures to restrict the construction of galleries, such as restoring gently sloping banks with abundant vegetation or introducing stony substrates, constitutes tangible elements that can be integrated into the development plan for a water body.



References

Crepin, E. (2021). *Élaboration d'un guide de bonnes pratiques pour la gestion des plantes aquatiques et des écrevisses envahissantes*, p. 223.

Holdich, D. M., Gydemo, R. and Rogers, W. D. (2017). *A review of possible methods for controlling nuisance populations of alien crayfish*. In: *Crayfish in Europe as alien species*, pp. 245-270.

Krieg, R., King, A. and Zenker, A. (2020). *Measures to Control Invasive Crayfish Species in Switzerland: A Success Story?*. *Frontiers in Environmental Science*, 8, 1-9.

Lemmers, P. et al. (2022). *Limiting burrowing activity and overland dispersal of the invasive alien red swamp crayfish *Procambarus clarkii* by sophisticated design of watercourses*. *Ecological Engineering*, 185, 1-8.

Poulet, N. (2014). *Les méthodes de contrôle des populations d'écrevisses invasives*, p. 13.

Fig 15. Natural banks are less likely to be invaded by crayfish species Photo: Arnaud Monty

The impact of management actions on ecosystem services

While the adverse effects of IAS are well-known and provide strong incentives for implementing management actions, the impacts of these management actions on ecosystems and the services they provide are less considered. The matrices are the result of expert assessments of the evolution of relevant ecosystem services (ES) from a highly invaded situation towards a managed situation. ES evolution is considered over 2 given periods of time: 1 year and 5 years after the initiation of management.




Fig 16. Representation of the survey process

Each matrix displays the average impact scores of management methods on ecosystem services. These scores have been associated to colours to facilitate the visualization of the impacts of every method on every relevant ecosystem service. Green indicates a significant improvement in the ecosystem services (ES) due to management, orange represents no or minimal effect, and red signifies a negative impact of the method on the ES.




Fig 17. Matrix displaying the impact of management methods for crayfish species on ecosystem services after 1 year




Fig 18. Matrix displaying the impact of management methods for crayfish species on ecosystem services after 5 years

LIFE RIPARIAS

Reaching Integrated and Prompt Action in Response to Invasive Alien Species

Citation

Monty, A., Latli A., Vermeersch, X. and Patinet, M. (2024). Management fact sheet - *Cherax destructor*. LIFE RIPARIAS project, 18p.

© LIFE RIPARIAS 2024. Licence CC-BY-NC. Licensed work, commercial use of this fact sheet is not permitted.

© Photo credits: This management fact sheet is extracted from the guide "Invasive Alien Crayfish Species - Identification and Best Management Practices." For a comprehensive list of photo credits, please consult the [full management guide](#).

www.riparias.be

This project is co-funded by the European Union. Views and opinions expressed are however those of the authors only and do not necessarily reflect those of the European Union or CINEA. Neither the European Union nor the granting authority can be held responsible for them.

RIPARIAS

Reaching Integrated and Prompt Action
in Response to Invasive Alien Species

